




1-800-524-5427

sales@icswaco.com

www.icswaco.com

P.O. Box 21056 Waco, TX 76702-1056



	
<h2 style="text-align: center;">Aminofen</h2>	
<p style="text-align: center;">SKU: 1625303</p>	
<p>Aspirin-free relief from pain and discomfort associated with colds, arthritis, headache, muscle ache, menstrual cramps and fever</p> <ul style="list-style-type: none"> • Unit dose, tamper-evident packets • PACKAGING IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN <p>Each tablet contains: Acetaminophen 325 mg</p>	
Pack	2/PACK, 250 PACKS/BOX
Weight	0.7687 lbs.
Ingredients	<p>Active ingredient (in each tablet): Acetaminophen 325 mg Purpose: Pain reliever/fever reducer</p> <p>Inactive ingredients* corn starch, hypromellose, maltodextrin*, microcrystalline cellulose*, polyethylene glycol, povidone*, pregelatinized starch*, sodium starch glycolate*, stearic acid, titanium dioxide*</p> <p>*may contain</p>
Uses	<p>For the temporary relief of minor aches and pains associated with\</p> <ul style="list-style-type: none"> • headache • muscular aches • minor arthritis pain • common cold • toothache • menstrual cramps <p>For the reduction of fever.</p>
Directions	<ul style="list-style-type: none"> • do not use more than directed <p>Adults and children: (12 years and older) Take 2 tablets every 4 to 6 hours while symptoms last. Do not take more than 10 tablets in 24 hours, unless directed by a doctor.</p> <p>Children under 12 years: Do not give to children under 12 years of age.</p>
Other Information	<ul style="list-style-type: none"> • store at room temperature 59°-86°F (15°-30°C) • tamper-evident sealed packets • do not use any opened or torn packets



1-800-524-5427

sales@icswaco.com

www.icswaco.com

P.O. Box 21056 Waco, TX 76702-1056



Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.