




1-800-524-5427

sales@icswaco.com

www.icswaco.com

P.O. Box 21056 Waco, TX 76702-1056



	<h2 style="text-align: center;">Medi-First Antacid</h2> <p style="text-align: center;">SKU: 80233</p> <p>Aluminum-free calcium supplement for heartburn, sour stomach, acid indigestion and upset stomach. Mint flavor.</p> <ul style="list-style-type: none"> • Great tasting, chewable tablets • Unit dose, tamper-evident packets • Spanish labeling where possible <p>Each tablet contains: Calcium Carbonate 420 mg</p>
<p>Pack</p>	<p>2/PACK, 50 PACKS/BOX</p>
<p>Weight</p>	<p>0.1937 lbs.</p>
<p>Ingredients</p>	<p>Active ingredient (in each tablet): Calcium Carbonate 420 mg Purpose: Antacid</p> <p>Inactive ingredients* aspartame*, croscarmellose sodium*, gum acacia*, magnesium stearate, maltodextrin, mineral oil*, mint flavor, sorbitol*, sucrose* <i>*may contain</i></p>
<p>Uses</p>	<p>For the relief of the following symptoms</p> <ul style="list-style-type: none"> • acid indigestion • sour stomach • heartburn • upset stomach associated with these symptoms
<p>Directions</p>	<ul style="list-style-type: none"> • do not use more than directed • chew or crush tablets completely before swallowing • do not swallow tablets whole <p>Adults and children: (12 years and older) Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not exceed 19 tablets in 24 hours. Children under 12 years: Do not give to children under 12 years of age.</p>
<p>Other Information</p>	<ul style="list-style-type: none"> • Phenylketonurics: may contain phenylalanine 1.5 mg per tablet • each tablet contains 168 mg of elemental calcium • store at room temperature 59°-86°F (15°-30°C) in a dry place • tamper-evident sealed packets • do not use any opened or torn packets



1-800-524-5427

sales@icswaco.com

www.icswaco.com

P.O. Box 21056 Waco, TX 76702-1056



Warnings

Do not use

- the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician, or take more than 19 tablets in a 24 hour period.

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

- symptoms last more than 2 weeks

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.